

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

Acids , Bases and Salts in Medical Interests

2nd lecture of Medical chemistry
For college of Medicine Students
by
Dr. Salih Mahdi Salman

What is an ACID?

- pH less than 7
- Neutralizes bases
- Forms H⁺ ions in solution
- Corrosive-reacts with most metals to form hydrogen gas
- Good conductors of electricity
- Acids generate ions



Weak & Strong Acids

- **Weak Acids** do not ionize completely: Acetic, Boric, Nitrous, Phosphoric, Sulfurous.
- **Strong Acids** ionize completely: Hydrochloric, Nitric; Sulfuric Hydriodic.

What is a BASE?

- pH greater than 7
- Feels slippery
- Dissolves fats and oils
- Usually forms OH^- ions in solution
- Neutralizes acids



Weak & Strong Bases

- **Weak Bases** do not ionize completely : ammonia, potassium carbonate, sodium carbonate.
- **Strong Bases** ionize completely : sodium hydroxide, sodium phosphate, barium hydroxide, calcium hydroxide

Definitions of acids and bases

Arrhenius

acid: generates $[H^+]$ in solution


base: generates $[OH^-]$ in solution



Bronsted-Lowry:

acid: anything that donates a $[H^+]$ (proton donor)

base: anything that accepts a $[H^+]$ (proton acceptor)




Lewis:

acid: accepts an electron pair


base: donates an electron pair

The advantage of this theory is that many reactions can be considered acid-base reactions because they do not have to occur in solution.



pH Scale

- The strength of an acid or base in a solution is measured on a scale called a pH scale or .
- The pH scale is a measure of the hydrogen ion concentration.




What is a SALT?



- A salt is a neutral substance produced from the reaction of an acid and a base.
- Composed of the negative ion of an acid and the positive ion of a base.
- One of the products of a Neutralization Reaction
Examples: KCl , MgSO_4 , Na_3PO_4

Source of acids and bases in the body


1. Inorganic acids produced during the breakdown of nutrients.



2. Organic acids resulting from intermediary metabolism.


- Fatty acids are produced during fat metabolism
 - Lactic acid is produced by muscles during heavy exercise.
- These acids partially dissociate to yield free H⁺

3. Carbonic acid formation



The Body and pH

- Homeostasis of pH is tightly controlled
- Extracellular fluid = 7.4
- Blood = 7.35 – 7.45
- < 6.8 or > 8.0 death occurs
- **Acidosis** (acidemia) below 7.35
- **Alkalosis** (alkalemia) above 7.45



Effects of fluctuations in hydrogen-ion concentration on body chemistry

The prominent consequences of fluctuations in $[H^+]$ include the following:

1. Changes in excitability of nerve and muscle cells.

- The major clinical effect of **increased $[H^+]$ (acidosis)** is **depression of the central nervous system.**
- The major clinical effect of **decreased $[H^+]$ (alkalosis)** is **over excitability of the nervous system**

2. Hydrogen-ion concentration exerts a marked influence on enzyme activity.

- Since enzymes are proteins, a shift in the body acid-base balance disturbs the normal pattern of metabolic activity catalyzed by these enzymes
- **Some cellular chemical reactions are accelerated; others are depressed.**

3. Changes in $[H^+]$ influence K^+ levels in the body.

- If more H^+ than normal is eliminated by the kidneys the body fluids become acidotic, less K^+ than usual can be excreted.
- **The resultant K^+ retention can affect cardiac function.**

Acid-base balance

- Hydrogen-ion generation therefore normally goes on continuously as a result of on going metabolic activities.
- **But**, in certain states, additional acids may be produced that further contribute to the total body pool of H^+ .

For example


- In diabetes mellitus, large quantities of keto acids may be produced as a result of abnormal fat metabolism.
- Some types of acid-producing medications may also add to the total load of H^+ that must be handled by the body.
- **Thus**, input of H^+ is unceasing, highly variable, and essentially unregulated.

How can keep the balance of the acid in the body?



Three lines of defense against changes in $[H^+]$ operate to maintain the $[H^+]$ of body fluids at a nearly constant level despite unregulated input:

(1) The chemical buffer systems.



(2) The respiratory mechanism of pH control.




(3) The renal mechanism of pH control.



1. The chemical buffer systems of the body

There are three major buffer system the pH of the body

a. Carbonate buffer



Excess acid (H_3O^+) in the body is neutralized by HCO_3^-



Equilibrium shifts left

Excess base (OH^-) reacts with the carbonic acid (H_2CO_3)



Equilibrium shifts right

pH of the blood buffer

The concentrations of H_2CO_3 and HCO_3^- in the blood are 0.0024M and 0.024M respectively

$\text{H}_2\text{CO}_3 / \text{HCO}_3^- = 1/10$ is needed to maintain the normal blood pH (7.35 – 7.45)

$$K_a = \frac{[\text{H}_3\text{O}^+][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]}$$


$$[\text{H}_3\text{O}^+] = K_a \frac{[\text{H}_2\text{CO}_3]}{[\text{HCO}_3^-]}$$

$$= 4.3 \times 10^{-7} \times \frac{0.0024}{0.024} = 4.3 \times 10^{-7} \times 0.10 = 4.3 \times 10^{-8}$$

$$\text{pH} = -\log(4.3 \times 10^{-8}) = 7.37$$

b. phosphate buffer


The phosphate buffer system ($\text{HPO}_4^{2-}/\text{H}_2\text{PO}_4^-$) plays a role in plasma and erythrocytes.



Any acid reacts with monohydrogen phosphate to form dihydrogen phosphate



The base is neutralized by dihydrogen phosphate



c. Proteins buffer

Proteins contain – COO⁻ groups, which, like acetate ions (CH₃COO⁻), can act as proton acceptors.

Proteins also contain – NH₃⁺ groups, which, like ammonium ions (NH₄⁺), can donate protons.

If acid comes into blood, hydronium ions can be neutralized by the – COO⁻ groups



If base is added, it can be neutralized by the – NH₃⁺ groups



2. The respiratory mechanism of pH control.

- Amount of blood carbon dioxide directly relates to amount of carbonic acid and therefore to concentration of H^+
- With increased respirations, less carbon dioxide remains in blood, hence less carbonic acid and fewer H^+
- With decreased respirations, more carbon dioxide remains in blood, hence more carbonic acid and more H^+ .


Carbon dioxide levels and pH affect respiratory centers

Hypoventilation increases blood carbon dioxide levels

Hyperventilation decreases blood carbon dioxide levels

Respiratory Acidosis

- **Carbonic acid excess** caused by blood levels of CO_2 above 45 mm Hg.
- **Hypercapnia** – high levels of CO_2 in blood



Compensation for Respiratory Acidosis

- **The body response to acid-base imbalance is called compensation**
- **Kidneys eliminate hydrogen ion and retain bicarbonate ion**


Treatment of Respiratory Acidosis

Restore ventilation

IV lactate solution



a) Metabolic balance before onset of acidosis



H_2CO_3 : Carbonic acid
 HCO_3^- : Bicarbonate ion
 ($Na^+ \bullet HCO_3^-$)
 ($K^+ \bullet HCO_3^-$)
 ($Mg^{++} \bullet HCO_3^-$)
 ($Ca^{++} \bullet HCO_3^-$)


b) Respiratory acidosis




Primary change
 pH — decreases
 PCO_2 — increases
 HCO_3^- — no change

Breathing is suppressed, holding CO_2 in body

c) Body's compensation



d) Therapy required to restore metabolic balance



From Thibodeau GA, Patton KT: *Anatomy & physiology*, ed 5, St Louis, 2003, Mosby. Mosby items and derived items copyright © 2004, 2000 by Mosby, Inc.

Respiratory Alkalosis


- Carbonic acid deficit
- $p\text{CO}_2$ less than 35 mm Hg (hypocapnea)
- Most common acid-base imbalance
- Primary cause is hyperventilation

Compensation of Respiratory Alkalosis


- Kidneys conserve hydrogen ion
- Excrete bicarbonate ion

Treatment of Respiratory Alkalosis

- Breathe into a paper bag
- IV Chloride containing solution
- – Cl^- ions replace lost bicarbonate ions



a) Metabolic balance before onset of alkalosis




H_2CO_3 : Carbonic acid
 HCO_3^- : Bicarbonate ion
 ($Na^+ \bullet HCO_3^-$)
 ($K^+ \bullet HCO_3^-$)
 ($Mg^{++} \bullet HCO_3^-$)
 ($Ca^{++} \bullet HCO_3^-$)

b) Respiratory alkalosis





Hyperactive breathing "blows off" CO₂




Primary change
 pH — increases
 PCO₂ — decreases
 HCO₃⁻ — no change

c) Body's compensation



d) Therapy required to restore metabolic balance



From Thibodeau GA, Patton KT: *Anatomy & physiology*, ed 5, St Louis, 2003, Mosby.
 Mosby items and derived items copyright © 2004, 2000 by Mosby, Inc.

3.The renal mechanism of pH control.

- Can eliminate large amounts of acid
- Can also excrete base
- Can conserve and produce bicarbonate ions
- Most effective regulator of pH
- If kidneys fail, pH balance fails

Metabolic Acidosis

- **Bicarbonate deficit** - blood concentrations of bicarbonate drop below 22mEq/L
- **Causes:**
 - Loss of bicarbonate through diarrhea or renal dysfunction
 - Accumulation of acids (lactic acid or ketones)
 - Failure of kidneys to excrete H⁺

Symptoms of Metabolic Acidosis


- Headache, lethargy
- Nausea, vomiting, diarrhea
- Coma
- Death

Compensation for Metabolic Acidosis

- Increased ventilation
- Renal excretion of hydrogen ions if possible
- K^+ exchanges with excess H^+ in ECF (H^+ into cells, K^+ out of cells)

Treatment of Metabolic Acidosis

- IV lactate solution



Metabolic Alkalosis

- **Bicarbonate excess** - concentration in blood is greater than 26 mEq/L
- **Causes:**
 - Excess vomiting = loss of stomach acid
 - Excessive use of alkaline drugs
 - Certain diuretics
 - Endocrine disorders
 - Heavy ingestion of antacids
 - Severe dehydration


Compensation for Metabolic Alkalosis

- Alkalosis most commonly occurs with renal dysfunction, so can't count on kidneys
- Respiratory compensation difficult – hypoventilation limited by hypoxia

Treatment of Metabolic Alkalosis


- Electrolytes to replace those lost
- IV chloride containing solution

a) Metabolic balance before onset of alkalosis




H_2CO_3 : Carbonic acid
 HCO_3^- : Bicarbonate ion
 ($\text{Na}^+ \bullet \text{HCO}_3^-$)
 ($\text{K}^+ \bullet \text{HCO}_3^-$)
 ($\text{Mg}^{++} \bullet \text{HCO}_3^-$)
 ($\text{Ca}^{++} \bullet \text{HCO}_3^-$)

b) Metabolic alkalosis
 HCO_3^- increases because of loss of chloride ions or excess ingestion of sodium bicarbonate




Primary change
 pH — increases
 PCO_2 — no change
 HCO_3^- — increases

c) Body's compensation



d) Therapy required to restore metabolic balance



Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

**First line of
defense against
pH shift**

**Chemical
buffer system**

**Bicarbonate
buffer system**

**Phosphate
buffer system**

**Protein
buffer system**

**Second line of
defense against
pH shift**

**Physiological
buffers**

**Respiratory
mechanism
(CO₂ excretion)**

**Renal
mechanism
(H⁺ excretion)**

Rates of correction

- Buffers function almost instantaneously
- Respiratory mechanisms take several minutes to hours
- Renal mechanisms may take several hours to days

Diagnosis of Acid-Base Imbalances

1. Note whether the pH is low (acidosis) or high (alkalosis)
2. Decide which value, is outside the normal range and could be the

cause of the problem $p\text{CO}_2$ or HCO_3^- .

If the cause is a change in $p\text{CO}_2$, the problem is respiratory.

If the cause is HCO_3^- the problem is metabolic.

3. Look at the value that doesn't correspond to the observed pH change whether its **inside** or outside the normal range.

If it is inside the normal range, there is no compensation occurring.

If it is outside the normal range, the body is partially compensating for the problem.

Example:- A patient is in intensive care because he suffered a severe myocardial infarction 3 days ago. The lab reports the following values from an arterial blood sample:

pH 7.3

$\text{HCO}_3^- = 20 \text{ mEq / L (22 - 26)}$

$\text{pCO}_2 = 32 \text{ mm Hg (35 - 45)}$

**Diagnosis : Metabolic acidosis
With compensation**

A magical night landscape featuring a full moon in the upper left, a vibrant aurora borealis in shades of green and blue across the sky, and a field of glowing purple flowers in the foreground. The flowers have a bright, starry glow emanating from their centers. The background shows dark, silhouetted mountains under a starry night sky.

Thank you for your attention