

Fat-soluble vitamins

- **A, D, E and K**

Vitamin (A)

- Vitamin A was the first fat-soluble vitamin to be discovered and has been known chemically since 1927
- Chemistry:
- The term 'vitamins A' is reserved to design any substance or mixture of substances that possesses activity similar to vit. A. **Retinol** is the most commonly known and abundant of all **naturally** occurring vit.A compounds.
- Food Sources:
- Performed vit.A is available only in foods of animal origin in which the animal has converted the precursor into active vitamin. Rich sources of the vitamin include: liver, kidney, and animal fats including these found in milk, butter, egg yolk and fatty fish. [retinol—natural source of vit.A].
- Plants food are rich in **carotenoids** which are chemically related to vit.A. They are known as the precursors (or provitamin A) that are converted in the body by an enzyme-catalyzed reaction to active vit.A. The major dietary sources of provitamin A are the yellow and green vegetables and fruits [carrots, sweet potatoes, apricots, spinach]. The carotenes are useful nutritionally only in so far as they may be converted to retinol.[pumpkin, tomato, peach, water melon]

Function of vitamin A

1. Vision: the light receptors of the eye are the **rod** and **cone** cells of the retina, both contain light –sensitive pigments that require vit A for their formation and proper functioning.
2. Maintenance of epithelial cells: retinol deficiency has a deleterious effect on epithelial structure in general. Epithelial cells are found in the linings of all openings in the interior of the body as well as in glands and ducts. They also form the outer protective layer of the skin. Most epithelial cells secrete mucus, but in vitamin A deficiency there is reduction in mucus secreting cell.
3. Bone development: vitamin A is required for normal bone growth. In a deficiency, bones fail to grow in length, probably because the remodeling process that is an essential phase of bone growth is poorly controlled. The skull and spinal column do not continue to enlarge to accommodate the growing nervous system. Vitamin A is also necessary for enamel forming epithelial cells in the development of teeth antioxidant factor against cancer I.e. lung cancer.

Vitamin A requirement

1. Influencing factors a number of factors modify the needs such as:

- A.** The amount stored in the liver
- B.** The form in which it is taken (carotene or vit A), as only approximately (1/3) of beta carotene is absorbed in to intestinal wall and then only (1/2) of this is converted in to retinol. The absorption of both retinol and carotene is facilitated by fats in the diet and bile salt in the duodenum.
- C.** Illness
- D.** Gastrointestinal or hepatic defects.

2. Units: the units are presently expressed as retinol equivalents (R.E). It takes in to account the variations in absorption and conversion of different precursors in to vit A. One R.E is equal to 1µg of retinol, 6 µg of β-carotene and 12 µg of other carotene precursors. Which vit.A is expressed in international Units (I.U), 1 R.E. =3.3 I.U. of retinol=10 I.U. of carotene.

3. RDA:

The requirement of vit. A is proportional to body weight the RDA for adult males is 1000 R.E; the RDA for women is lower, 800 R.E, during pregnancy , 1000 R.E. are recommended and during lactation 1200 R.E., children need 600-1000 R.E. daily, with the amount increasing from infancy to 14 years of age.

Vitamin A Deficiency

- **Primary vit.A deficiency:** - generally results from inadequate intake in the diet, a low intake of protein and fat needed for the absorption and utilization of vit. A may aggravate the problem. The liver can store quantities of vit. A sufficient for several months and therefore deficiency only becomes apparent when inadequate intake has persisted for so long that the liver stores are too depleted to supply the body's requirement.
- **Secondary vit. A deficiency** occurs due to:-
 1. Poor absorption due to lack of bile or defective absorbing surface
 2. Inadequate conversion of carotene because of liver disease (liver cirrhosis) or intestinal disease

Young children because of their requirements are particularly at risk of vit A deficiency, as vit A deficiency and its ocular clinical manifestation xerophthalmia is found mainly among the poor of the developing countries. Vit A deficiency is one of the seven most common causes of blindness. The WHO estimates that 250 thousand children become blind every year from vit A deficiency. Clinical features :earliest symptom is night blindness followed by degenerative changes in the retina.

Xerophthalmia Classification

- **XN Night blindness**
- **X1A Conjunctival xerosis(conjunctiva becomes dry and thick)**
- **X1B Bitot's spot—small foamy, superficial plaque on the temporal conjunctiva.**
- **X2 Corneal xerosis as the cornea becomes dry and lacks luster, later it may look bluish and milky.**
- **X3A Corneal ulceration with xerosis.**
- **X3B Keratomalacia.**
- **XF Xerophthalmia Fundus, softening of the cornea that leads to permanent scarring and deformation of the cornea, leading to blindness.**
- **XS Scare.**

Vitamin A deficiency is considered to be a public health problem if, among children aged 0.5 years, at least if;
1% have night blindness or
2% have Bitot's Spots or
0.01% have corneal dryness, ulcers or keratomalacia or
0.05% have corneal ulcers.

Vitamin A deficiency is associated with an increased risk of transmitting HIV from mother to infant in the breast milk. Severe forms of the disease are often associated with other co-existing factors that may affect both levels of and demand for, retinol binding protein. These include energy nutrient malnutrition, and infections. An association between measles and xerophthalmia has been reported. When the measles virus infects the conjunctiva and cornea, the epithelia are damaged; if there is insufficient vit.A or retinol binding protein available for epithelial regeneration. Xerophthalmia may develop. Also skin changes which include dryness and hyperkeratosis are also sign of vit. A deficiency.

Seasonal Pattern:

Xerophthalmia prevalence is greatest during the hot, dry months when green leafy vegetables are least prevalent and the incidence of infectious diseases are high.

Treatment

- **Immediately on diagnosis (60 mg) retinol (200,000 I.U) should be given orally, or if there is vomiting or severe diarrhea (55 mg) retinol by I.M injection. The corneal lesions clear up within 48-72 hours.**
- **The vitamin preparation in a dose of 8-10 mg/day should be given for a month. Patients with ocular complication should be referred to the ophthalmologist.**

Prevention

- **The control of vitamin A deficiency can only be achieved through an improvement in the population's dietary intake of retinol or beta carotene, this may require a change in food habits. Pregnant woman should be advised to eat dark green leafy vegetables. This helps to build up stores of retinol in the fetal liver and should be educated about the importance of given such vegetables to their babies.**
- **A more specific short term measure is the oral administration of single high dose capsule containing 200.000 I.U of vit. A to-- at risk children.**
- **In Iraq, we provide a single dose of vit. A (50.000) I.U to young children at the age of 9 months with the measles vaccine, at the age of 18 months another dose of vit.A (100.000) I.U, and another prophylactic dose at the first year of primary school, also a single dose of (200.000) I.U usually provided to lactating mothers after delivery.**

Vitamin D

Vit D is required for normal metabolism of calcium and phosphorus and for bone formation. It enhances the absorption of these minerals from the gut, their mobilization from bone and the reabsorption of phosphorus by the kidney.

All forms of vit D are soluble in fat but not in water.

Dietary Sources of vit. D

Few natural food sources of vit.D exist. The main food sources are milk, butter, cheese, egg yolk and fish liver oils. In general foods are low sources of vit.D, as human milk contain 2-100 I.U/L, cow milk 3-50 I.U/L, egg yolk 100-500 I.U/100gram, 20% of egg is egg yolk, therefore 7 eggs are needed to compensate for this quantity.

Therefore the **main source is the skin synthesis; smoke prevented the Ultra Violet Radiation in crowded urban areas.**

Vitamin D Requirement

- Influencing factors: difficulties exist in setting requirements for vit.D. Variables arise from many sources:-
 1. A limited number of food sources available.
 2. Lack of knowledge of precise body needs.
 3. Differing degrees of skin synthesis by irradiation.
 4. Need; vary between winter and summer.
 5. A persons way of living and working determines the degree of **sunlight exposure**. Elderly people who do not go out of doors may need supplementary vit.D. Growth demands in children and in pregnancy and lactation necessitate increase in take.
- In general RDA standard is 400 I.U (10µg).

Vitamin Deficiency

- Vitamin D deficiency causes **Rickets** in the growing child and **Osteomalacia** in adult.
- **Aetiology of vitamin Deficiency:**
These disorders resulted from various aetiological factors which all results in deficient calcium and phosphorous metabolism.
 1. Deficient **diet** intake of vitamin D.
 2. Inadequate exposure to **sunlight**.
 3. **Malabsorption** diseases such as celiac syndrome, colitis and crohn disease.
 4. Anticonvulsant **drug** therapy which transform vit.D into inactive form, or it effect the transfer of vit. D into an active form.
 5. Congenital defect of some **enzymes**.
 6. Defective tubular reabsorption which lead to loss of phosphorous and to less degree calcium.
 7. Chronic **renal** insufficiency which is called Osteodystrophy.

Treatment

- Oral administration of vitamin D in doses of **1500-5000 I.U daily** brings about rapid improvement in vast majority of cases. The dose may be given orally or as I.M injection. After complete healing of the lesion, vit. D should be given in dose of 400 I.U daily for preventing recurrence. Children should be encourage to get exposure to **sun** in moderate amounts.

Prevention

- **Supplementation of the infant food with vit.D is required, as fortified milk or baby food with 100 I.U/L, but there is the danger of vit.D intoxication can occurred from excessively fortified food, or vit.D can be given by drops for daily supplement of 400 I.U, and it is shown that it is better absorbed than fortify milk.**

Thank you

