

Cyanotic Congenital Heart Disease Lesions Associated with Increased Pulmonary Blood Flow

D-Transposition of the Great Arteries

Transposition of the great vessels, a common cyanotic congenital anomaly, accounts for $\approx 5\%$ of all congenital heart disease. In this anomaly, the systemic veins return normally to the right atrium and the pulmonary veins return to the left atrium. The connections between the atria and ventricles are also normal (atrioventricular concordance). The aorta arises from the right ventricle and the pulmonary artery from the left ventricle. Desaturated blood returning from the body to the right side of the heart goes inappropriately right out the aorta and back to the body again, whereas oxygenated pulmonary venous blood returning to the left side of the heart is returned directly to the lungs. The systemic and pulmonary circulations consist of two parallel circuits. Survival in these newborns is provided by the foramen ovale and the ductus arteriosus, which permit some mixture of oxygenated and deoxygenated blood. About 50% of patients with TGA also have a ventricular septal defect (VSD), which provides for better mixing. The clinical findings and hemodynamics vary in relation to the presence or absence of associated defects. TGA is more common in infants of diabetic mothers and in males (3 : 1). d-TGA, especially when accompanied by other cardiac defects such as pulmonic stenosis or right aortic arch, can be associated with deletion of chromosome 22q11.2 (DiGeorge syndrome). Before the modern era of corrective or palliative surgery, mortality was $>90\%$ in the 1st yr of life.

D-Transposition of the Great Arteries with Intact Ventricular Septum

D-TGA with an intact ventricular septum is also referred to as simple TGA or isolated TGA. Before birth, oxygenation of the fetus is nearly normal, but after birth, once the ductus begins to close, the minimal mixing of systemic and pulmonary blood via the patent foramen ovale is usually insufficient and severe hypoxemia ensues, generally within the 1st few days of life.

CLINICAL MANIFESTATIONS

Cyanosis and tachypnea are most often recognized within the 1st hrs or days of life. Untreated, the vast majority of these infants would not survive the

neonatal period. Hypoxemia is usually severe, but heart failure is less common. This condition is a medical emergency, and only early diagnosis and appropriate intervention can avert the development of prolonged severe hypoxemia and acidosis, which lead to death. The 2nd heart sound is usually single and loud, although occasionally it may be split. Murmurs may be absent, or a soft systolic ejection murmur may be noted at the mid left sternal border.

DIAGNOSIS

The electrocardiogram shows the normal neonatal right-sided dominant pattern. Roentgenograms of the chest may show mild cardiomegaly, a narrow mediastinum (hence an egg-shaped heart), and normal to increased pulmonary blood flow. Arterial PO₂ is low and does not rise appreciably after the patient breathes 100% oxygen (hyperoxia test), although this test may not be totally reliable. Echocardiography confirms the transposed ventricular-arterial connections. The degree of mixing is assessed by pulsed and color Doppler examination. Cardiac catheterization is occasionally performed in patients for whom noninvasive imaging is diagnostically inconclusive or in patients who require emergency balloon atrial septostomy.

TREATMENT

When transposition is suspected, an infusion of prostaglandin E₁ should be initiated immediately to maintain patency of the ductus arteriosus and improve oxygenation (dosage, 0.01–0.20 µg/kg/min). Because of the risk of apnea associated with prostaglandin infusion, an individual skilled in neonatal endotracheal intubation should be available. Hypothermia intensifies the metabolic acidosis resulting from hypoxemia, and thus the patient should be kept warm. Infants who remain severely hypoxic or acidotic despite prostaglandin infusion should undergo Rashkind balloon atrial septostomy. A Rashkind atrial septostomy is usually performed in all patients in whom any significant delay in surgery is necessary.

The arterial switch (Jantene) procedure is the surgical treatment of choice for neonates with d-TGA and an intact ventricular septum and is usually performed within the 1st 2 wk of life. The arterial switch procedure has a

survival rate of 90–95% for uncomplicated d-TGA. It restores the normal physiologic relationships of systemic and pulmonary arterial blood flow and eliminates the long-term complications of the atrial switch procedure.

Transposition of the Great Arteries with Ventricular Septal Defect

If the VSD associated with TGA is small, the clinical manifestations, laboratory findings, and treatment are similar to those described previously for transposition with an intact ventricular septum. A harsh systolic murmur is audible at the lower left sternal border and results from flow through the defect. When the VSD is large and not restrictive to ventricular ejection, significant mixing of oxygenated and deoxygenated blood usually occurs and clinical manifestations of cardiac failure are seen. The murmur is holosystolic and generally indistinguishable from that produced by a large VSD in patients with normally related great arteries. The heart is usually significantly enlarged.

In equivocal cases, the diagnosis can be confirmed by cardiac catheterization. At the time of cardiac catheterization, Rashkind balloon atrial septostomy may be performed to decompress the left atrium, even when adequate mixing is occurring at the ventricular level. Surgical treatment is advised soon after diagnosis, usually within the 1st months of life, because heart failure and failure to thrive are difficult to manage and pulmonary vascular disease can develop unusually rapidly. Management includes diuretics and, possibly, digitalis to lessen the symptoms of heart failure while awaiting surgical repair.

L-Transposition of the Great Arteries (Corrected Transposition)

In l-transposition, the atrioventricular relationships are discordant, with the right atrium connected to the left ventricle and the left atrium to the right ventricle (ventricular inversion). The great arteries are also transposed, with the aorta arising from the right ventricle and the pulmonary artery from the left. Desaturated systemic venous blood is returned to a normally positioned right atrium, from which it passes through a bicuspid atrioventricular (mitral) valve into a right-sided ventricle that has the architecture and smooth wall morphologic features of the normal left ventricle.

Total Anomalous Pulmonary Venous Return PATHOPHYSIOLOGY

Abnormal development of the pulmonary veins may result in either partial or complete anomalous drainage into the systemic venous circulation. Partial anomalous pulmonary venous return is usually an a cyanotic lesion.

In TAPVR, the heart has no direct pulmonary venous connection into the left atrium. The pulmonary veins may drain above the diaphragm into the right atrium directly, into the coronary sinus, or into the superior vena cava via a “vertical vein,” or they may drain below the diaphragm and join into a “descending vein” that enters into the inferior vena cava or one of its major tributaries, often via the ductus venosus.

All forms of TAPVR involve mixing of oxygenated and deoxygenated blood before or at the level of the right atrium (total mixing lesion). Right atrial blood either passes into the right ventricle and pulmonary artery or passes through an atrial septal defect (ASD) or patent foramen ovale into the left atrium. The manifestations of TAPVR depend on the presence or absence of obstruction of the venous channels.

CLINICAL MANIFESTATIONS

Two major clinical patterns of TAPVR are seen. Some are manifested in the neonatal period as severe obstruction to pulmonary venous return, most prevalent in the infracardiac group . Cyanosis and severe tachypnea are prominent, but murmurs may not be present These infants are severely ill and fail to respond to mechanical ventilation. Rapid diagnosis and surgical correction are necessary for survival. In contrast,those with mild or no obstruction to pulmonary venous return are usually characterized by the development of heart failure as the pulmonary vascular resistance falls, with mild to moderate degrees of desaturation.. Systolic murmurs are audible along the left sternal border.

DIAGNOSIS

The electrocardiogram demonstrates right ventricular hypertrophy, and the P waves are frequently tall and spiked) In neonates with marked pulmonary venous obstruction, the chest x-ray demonstrates a very dramatic perihilar pattern of pulmonary edema and a small heart. This appearance can sometimes be confused with primary pulmonary disease and the differential diagnosis includes persistent pulmonary hypertension of the newborn, respiratory distress syndrome, pneumonia (bacterial, meconium aspiration), pulmonary lymphangiectasia, and Roentgenograms are pathognomonic in

older children if the anomalous pulmonary veins enter the innominate vein and persistent left superior vena cava. A large supracardiac shadow can be seen, which together with the normal cardiac shadow forms a “snowman” appearance.

The echocardiogram demonstrates a large right ventricle and usually identifies the pattern of abnormal pulmonary venous connections. Cardiac catheterization shows that the oxygen saturation of blood in both atria, both ventricles, and the aorta is more or less similar, indicative of a total mixing lesion. MRI and CT may be alternative methods of confirming the diagnosis.

TREATMENT

Surgical correction of TAPVR is indicated during infancy. Before surgery, infants may be stabilized with prostaglandin E1 to dilate the ductus venosus and the ductus arteriosus, although with significant obstruction this is usually not effective. If surgery cannot be performed urgently.

Tetralogy of Fallot

Tetralogy of Fallot is one of the conotruncal family of heart lesions in which the primary defect is an anterior deviation of the infundibular septum (the muscular septum that separates the aortic and pulmonary outflows). The consequences of this deviation are the 4 components (1) obstruction to right ventricular outflow (pulmonary stenosis), (2) ventricular septal defect (VSD), (3) dextroposition of the aorta with override of the ventricular septum, and (4) right ventricular hypertrophy . Obstruction to pulmonary arterial blood flow is usually at both the right ventricular infundibulum (subpulmonic area) and the pulmonary valve. Complete obstruction of right ventricular outflow (pulmonary atresia with VSD) is classified as an extreme form of tetralogy of Fallot.

PATHOPHYSIOLOGY

More commonly, the subpulmonic or infundibular muscle, the crista supraventricularis, is hypertrophic, which contributes to the subvalvar stenosis and. When the right ventricular outflow tract is completely obstructed (pulmonary atresia. Pulmonary blood flow may be supplied by a patent ductus arteriosus (PDA) and by multiple major aortopulmonary

collateral arteries (MAPCAs) arising from the ascending and descending aorta.

The VSD is usually nonrestrictive and large, is located just below the aortic valve. The aortic arch is right sided in 20% of cases, and the aortic root is usually large and overrides the VSD to a varying degree.

When obstruction to right ventricular outflow is mild to moderate and a balanced shunt is present across the VSD, the patient may not be visibly cyanotic (acyanotic or “pink” tetralogy of Fallot). When obstruction is severe, cyanosis will be present from birth and worsen when the ductus begins to close.

CLINICAL MANIFESTATIONS

Often, cyanosis is not present at birth, but with increasing hypertrophy of the right ventricular infundibulum and patient growth, cyanosis occurs later in the 1st yr of life. Older children with long-standing cyanosis who have not undergone surgery may have dusky blue skin, gray sclerae with engorged blood vessels, and marked clubbing of the fingers and toes.

Characteristically, children assume a squatting position for the relief of dyspnea caused by physical effort; the child is usually able to resume physical activity within a few minutes. Paroxysmal hypercyanotic attacks (hypoxic, “blue,” or “tet” spells) are a particular problem during the 1st 2 yr of life. The spells occur most frequently in the morning on initially awakening or after episodes of vigorous crying. The spells may last from a few minutes to a few hours but are rarely fatal. Short episodes are followed by generalized weakness and sleep. Severe spells may progress to unconsciousness and, occasionally, to convulsions or hemiparesis. Infants who are only mildly cyanotic at rest are often more prone to the development of hypoxic spells because they have not acquired the homeostatic mechanisms to tolerate rapid lowering of arterial oxygen saturation, such as polycythemia.

Depending on the frequency and severity of hypercyanotic attacks, one or more of the following procedures should be instituted in sequence:-

(1) Placement of the infant on the abdomen in the knee-chest position while making certain that the infant's clothing is not constrictive.

(2) Administration of oxygen (although increasing inspired oxygen will not reverse cyanosis caused by intracardiac shunting).

(3) Injection of morphine subcutaneously in a dose not in excess of 0.2 mg/kg. Calming and holding the infant in a knee-chest position may abort progression of an early spell.

For spells that are resistant to this therapy, drugs that increase systemic vascular resistance, such as intravenous phenylephrine, can improve right ventricular outflow. β -Adrenergic blockade by the intravenous administration of propranolol is also useful. The murmur is caused by turbulence through the right ventricular outflow tract, it can actually become less prominent with severe obstruction, especially during a hypercyanotic spell.

DIAGNOSIS

The cardiac silhouette has been likened to that of a boot or wooden shoe. The electrocardiogram demonstrates right axis deviation and evidence of right ventricular hypertrophy. Cardiac catheterization demonstrates a systolic pressure in the right ventricle equal to systemic pressure.

COMPLICATIONS

Cerebral thromboses, are common in the presence of extreme Polycythemia and dehydration. Thromboses occur most often in patients younger than 2 yr. These patients may have iron deficiency anemia. Therapy consists of adequate hydration and supportive measures. Brain abscess is less common than cerebral vascular events and extremely rare when most patients are repaired at young ages. Patients with a brain abscess are usually older than 2 yr. Seizures may occur; localized neurologic signs depend on the site and size of the abscess and the presence of increased intracranial pressure. CT or MRI confirms the diagnosis. Antibiotic therapy may help keep the infection localized, but surgical drainage of the abscess is usually necessary.

Bacterial endocarditis may occur in the right ventricular infundibulum or on the pulmonic, aortic, or, rarely, tricuspid valves. Antibiotic prophylaxis is essential before and after dental and certain surgical procedures. Heart failure is not a usual feature in patients with the tetralogy of Fallot. It may occur in a young infant with “pink” or acyanotic tetralogy of Fallot. As the degree of pulmonary obstruction worsens with age, the symptoms of heart failure

resolve and eventually the patient experiences cyanosis, often by 6–12 mo of age.