

# Malignant tumors

د. محمد محمود حبش

# Classification

Cancers are classified by the type of cell that resembles the tumour and, therefore, the tissue presumed to be the origin of the tumour. These are the histology and the location, respectively. Examples of general categories include

**Carcinoma:** Malignant tumors derived from epithelial cells. This group represents the most common cancers, including the common forms of breast, prostate, lung and colon cancer

**Sarcoma:** Malignant tumors derived from connective tissue, or mesenchymal cells

**Lymphoma and leukemia:** Malignancies derived from hematopoietic cells

**Germ cell tumor:** Tumors derived from totipotent cells. In adults most often found in the testicle and ovary

**Blastic tumor or blastoma :** A tumor (usually malignant) which resembles an immature or embryonic tissue. Many of these tumors are most common in children

## Signs and symptoms

Symptoms of cancer metastasis depend on the location of the tumor

:Roughly, cancer symptoms can be divided into three groups

**Local symptoms:** unusual lumps or swelling (tumor), hemorrhage (bleeding), pain and/or ulceration. Compression of surrounding tissues may cause symptoms such as jaundice

**Symptoms of metastasis** (spreading): enlarged lymph nodes, cough and hemoptysis, hepatomegaly (enlarged liver), bone pain, fracture of affected bones and neurological symptoms. Although advanced cancer may cause pain, it is often not the first symptom

**Systemic symptoms:** weight loss, poor appetite, fatigue and cachexia (wasting), excessive sweating (night sweats), anemia and specific paraneoplastic phenomena, i.e. specific conditions that are due to an active cancer, such as thrombosis or hormonal changes

## Causes

Cancers are primarily an **environmental** disease with 90-95% of cases due to environmental factors and 5-10% due to **genetics**.

Most environmental causes, such as naturally occurring background **radiation**, are not modifiable or controllable

Common environmental factors that lead to cancer death include: **tobacco** (25-30% of deaths), **diet** and **obesity** (30-35%), **infections**(15-20%), **radiation**, **stress**, **lack of physical activity**, and **environmental pollutants**

## Chemicals

The incidence of lung cancer is highly correlated with smoking. Prolonged exposure to asbestos fibers is associated with .mesothelioma

*Alcohol is an example of a chemical carcinogen that is not a .mutagen*

Such chemicals may promote cancers through stimulating the rate of cell division. Faster rates of replication leaves less time for repair enzymes to repair damaged DNA during DNA replication, .increasing the likelihood of a mutation

## Radiation

Sources of ionizing radiation, such as radon gas, can cause cancer. Prolonged exposure to ultraviolet radiation from the sun can lead to melanoma and other skin malignancies.

Non-ionizing radio frequency radiation from mobile phones and other similar RF sources has also been proposed as a cause of cancer, but there is currently little established evidence of such a link

## **Infection**

Some cancers can be caused by infection. In humans, the viruses responsible for up to **20%** of human cancers worldwide. These include

, human papillomavirus (**cervical carcinoma**)

Epstein-Barr virus (**B-cell lymphoproliferative disease**, and **nasopharyngeal carcinoma**)

Kaposi's sarcoma herpesvirus (**Kaposi's Sarcoma**

hepatitis B and hepatitis C viruses (**hepatocellular carcinoma**), and

Human T-cell leukemia virus-1 (**T-cell leukemias**).

**Bacterial infection** may also increase the risk of cancer, as seen in **Helicobacter pylori induced gastric carcinoma**.

**HIV is associated with a number of malignancies, including Kaposi's sarcoma, non-Hodgkin's lymphoma, and HPV-associated malignancies such as anal cancer .and cervical cancer**



# Hereditary

Most forms of cancer are sporadic, meaning that there is no inherited cause of the cancer. There are, however, a number of recognized syndromes where there is an inherited predisposition to cancer, often due to a defect in a gene that protects against tumour formation. **Famous examples are**

- ?** certain inherited mutations in the **genes BRCA1 and BRCA2 are associated with an elevated risk of breast cancer and ovarian cancer**
- ?** tumors of various endocrine organs in multiple endocrine neoplasia(**MEN types 1, 2a, 2b**)
- ?** **Li-Fraumeni syndrome (various tumors such as osteosarcoma, breast cancer, soft tissue sarcoma, brain tumors) due to mutations of p53**
- ?** **Turcot syndrome(brain tumors and colonic polyposis)**
- ?** **Familial adenomatous polyposis an inherited mutation of the .APC gene that leads to early onset of colon carcinoma**

**❓ Hereditary nonpolyposis colorectal cancer(HNPCC, also known as Lynch syndrome) can include familial cases of colon cancer, uterine cancer, gastric cancer, and ovarian cancer, without a .preponderance of colon polyps**

**❓ Retinoblastoma, when occurring in young children, is due to a .hereditary mutation in the retinoblastoma gene**

**❓ Down syndrome patients, who have an extra chromosome 21, are known to develop malignancies such as leukemia and testicular cancer, though the reasons for this difference are not .well understood**

# Grading and staging

**Staging** is the process of assessing the extent of local and systemic spread of a malignant tumour or the identification of features that are risk factors for spread

**Grading** is the process of assessing the degree of differentiation of a malignant tumour

**:The objectives of staging and grading a tumour are**

;to plan appropriate (treatment) for the individual patient **\_1**

;to give an estimate of the prognosis **\_2**

to compare similar cases when assessing outcomes or designing **\_3**  
clinical trials

# Staging and grading methods

## Staging

The commonest system is the internationally agreed **TNM** classification. **It is not appropriate for leukemia lymphomas, or myeloma.** A four-stage classification (I, II, III, IV) is also often used and is compatible with TNM. Specific staging systems also exist for some tumour sites (e.g. Duke's stage in colorectal cancer)

***.Staging may be either radiological or pathological***

**Radiological** (often performed preoperatively) indicated by the prefix before the letter (e.g. rT3, rM1). If different radiological modalities are used separate prefixes can be used, e.g. for ultrasound (uT2). Radiological staging is used to plan treatment (e.g. neoadjuvant therapy, selection for surgery, planning of .surgery)

**Pathological** (performed on surgical specimens) indicated by the prefix before the letter (e.g. pT3, pN2, pM1). If there has been preoperative radiotherapy, the prefix is used to denote that the pathological stage may have been modified by this, e.g. ypT2. Pathological staging is used to plan adjuvant treatment .(chemotherapy or radiotherapy) and for informing prognosis

## An example of lung cancer staging

- Stage I (T1 NO, T2NO): **85%** 5y survival with surgery
- Stage II (T1 N1, T2N1, T3N0): **60%** 5y survival with  
.surgery
- Stage IIIa (T3 N1 or any N2): **20%** 5y survival with  
.surgery
- Stage IIIb (Any T4, any N3): **< 20%** 5y survival; no  
.benefit with surgery
- Stage IV(M1): **< 10%** 5y survival; no benefit with  
.surgery

**Other pathological features may be included with the .TNM system for some tumours, e.g**

☐ The presence of extra tumoural **vascular invasion** V0 or V1

☐ Presence of extra tumoural **lymphatic invasion** Ly0 or Ly1

☐ Presence of viable tumour cells at or within 1mm of the **surgical margin of excision** R0, R1 (microscopic), R2  
.(macroscopic)

## **Histological grading**

Gives a guide to the behaviour of a cancer by describing the .degree of differentiation of the tumour (e.g. breast cancer)

**.? Grade 1: represents the least malignant tumours**

**.? Grade 2: 25-50% of the cells are undifferentiated**

**.? Grade 3: 50-75% of the cells are undifferentiated**

**.? Grade 4: more than 75% of the cells are undifferentiated**



**Grading** predicts the aggressiveness of a malignant neoplasm by characterising its microscopic appearance taking into account

,The degree of differentiation .1

,Nuclear and cellular appearance .2

Architectural integrity and .3

.The proportion of active mitoses .4

**;Grade 1: well differentiated •**

**;Grade 2: moderately well differentiated •**

**.Grade 3: poorly differentiated •**

# **Factors affecting the prognosis of malignant tumour**

The prognosis of any tumour depends on four main features

**Extent of spread (1**

**Microscopic appearance (2**

**Anatomical situation (3**

**General condition of the patient (4**

## ***The extent of spread***

The extent of the tumour (it's staging) on clinical examination, operation and on studying the excised surgical specimen, is of great prognostic importance. Obviously, the clinical findings of palpable distant secondaries or gross fixation of the primary tumor .are serious

Similarly, the local invasiveness of the tumour at operation and .evidence of distant spread are of great significance

Finally histological study may reveal involvement of the nodes which had not detected clinically or microscopic extension of the growth at the edges of the resected specimen with consequent .worsening of the outlook for the patient

## ***Microscopic appearance of the tumour (histological differentiation)***

As general principle, the prognosis of tumor is related to its degree of histological differentiation (its grading) which is a spectrum between well differentiated and anaplastic

The spread of the tumour and its histological differentiation should be considered in conjunction with each other

A small tumour with no apparent spread at the time of operation may still have poor prognosis if it is highly anaplastic, whereas, an extensive tumour is not incompatible with long survival of the patient after operation if the microscopic examination reveals a high degree of differentiation


## ***Anatomic Situation***

The site of the tumour may preclude its adequate removal and thus seriously affect the prognosis e.g. tumour at the lower end of the esophagus may be easily removable whereas an exactly similar tumour situated behind the arch of the aorta may be technically inoperable

A brain tumour located in the frontal lobe may be resected whereas similar tumour in the brain stem will be not

## ***General condition of the patient***

A patient apparently curable from the point view of the local condition may be inoperable because of poor general health. For example, a gross congestive heart failure may convert what is technically operable carcinoma of the rectum into a hopeless anesthetic risk.





128600042







***Thank you***